



石蜡切片荧光探针原位杂交（FISH）实验报告

1.实验器材及试剂

1.1.实验器材

名称	厂家	型号
脱水机	DIAPATH	Donatello
载玻片	Servicebio	
盖玻片	江苏世泰实验器材有限公司	10212432C
病理切片机	上海莱卡仪器有限公司	RM2016
无酶离心管	Servicebio	EP-150-M
摇床(钟摆式)	Servicebio	TSY-B
涡旋混匀器	Servicebio	MV-100
移液枪	Dragon	KE0003087/KA0056573
免疫组化笔	Servicebio	G6100
冰箱	青岛海尔股份有限公司	BCD-192TGN
正置荧光显微镜	日本尼康	NIKON ECLIPSE CI
成像系统	日本尼康	NIKON DS-U3
恒温箱	LABOTERY	GSP-70
高压灭菌锅	松下健康医疗	MLS-3751L-PC
包埋机	武汉俊杰电子有限公司	JB-P5
冻台	武汉俊杰电子有限公司	JB-L5
组织摊片机	浙江省金华市科迪仪器设备有限公司	KD-P

1.2.主要实验试剂

试剂	厂家	货号	稀释比
DEPC	Amresco	E174	
4%多聚甲醛（DEPC 水）	Servicebio	G1113	
石蜡	Sakura		
无水乙醇	国药集团化学试剂有限公司	100092683	
环保型脱蜡透明液	Servicebio	G1128	
PBS	Servicebio	G0002	
20×SSC 洗脱液	Servicebio	G3016-4	
BSA	Servicebio	G5001	



蛋白酶 K	Servicebio	G1234
DAPI	Servicebio	G1012
抗荧光淬灭封片剂	Servicebio	G1401
杂交缓冲液	Servicebio	G3016-3

2. 石蜡切片荧光探针原位杂交实验步骤

- 2.1.组织固定：**组织取出洗净后立即放入固定液（DEPC 水配制）中固定 12h 以上。
- 2.2.脱水：**组织固定完成后经梯度酒精脱水后浸蜡，包埋。
- 2.3.切片：**石蜡经切片机切片，摊片机捞片，62°烤箱烤片 2h。
- 2.4.石蜡切片脱蜡至水：**依次将切片放入脱蜡透明液I15min-脱蜡透明液II15min-无水乙醇I5min-无水乙醇II5min,自然晾干，DEPC 水浸泡。
- 2.5.消化：**根据组织固定时间长短，切片于修复液中煮沸 10-15 分钟，自然冷却。后基因笔画圈，根据不同组织不同指标特性，滴加蛋白酶 K(20ug/ml) 37°消化____min。纯水冲洗后 PBS 洗 3 次×5min。
- 2.6.预杂交：**滴加预杂交液，37°C 孵育 1h。
- 2.7.杂交：**倾去预杂交液，滴加含探针____杂交液，浓度____，恒温箱____度杂交过夜。
- 2.8.杂交后洗涤：**洗去杂交液，2×SSC，37°C 洗 10min，1×SSC，37°C 洗 2×5min，0.5×SSC 室温洗 10min。若非特异杂交体较多，可以增加甲酰胺洗涤。
- 2.9.DAPI 复染核：**切片滴加 DAPI 染液，避光孵育 8min，冲洗后滴加抗荧光淬灭封片剂封片。
- 2.10.镜检拍照：**切片于尼康正置荧光显微镜下观察并采集图像。（紫外激发波长 330-380nm，发射波长 420nm,发蓝光；FAM(488)绿光激发波长 465-495nm，发射波长 515-555 nm，发绿光；CY3 红光激发波长 510-560，发射波长 590nm，发红光。）

3. 石蜡切片荧光探针原位杂交实验结果判读

DAPI 染出来的细胞核在紫外的激发下为蓝色，阳性表达为相应荧光素标记的荧光。FAM(488) 为绿光，cy3 为红光。mRNA 原位杂交显示结果理论为胞浆阳性，少数核阳性属正常。micRNA 与 lncRNA 不同指标表达定位不同。根据表达量不同荧光亮度有强弱。

注：上述涉及到的所有试剂，仪器等在 RNA 原位杂交实验时都需使用 DEPC 处理后的 RNase free 环境。

附表 1 探针信息



Paraffin-fluorescence Probe-FISH Protocol

1. Apparatus and Reagents

1.1 Apparatus

Name	Producer	Model
Dehydrator	WHJJ	JJ-12J
Paraffin embedding machine	WHJJ	JB-P5
Frozen flat	WHJJ	JB-L5
Pathologic microtome	Leica	RM2016
Water Bath-Slide	Kedee	KD-P
RNase-freeGlass microscope slides	Servicebio	
Micro-centrifuge	Servicebio	D1008E
Rocker	Servicebio	TSY-B
Vortex	Servicebio	MV-100
Pipettor	Dragon	KE0003087/KA0056573
Microscopy	Nikon	NIKON ECLIPSE CI
Imaging system	Nikon	NIKON DS-U3
Liquid Blocker PAP Pen	Servicebio	G6100
Refrigerator	HAIER	BCD-192TGN
Incubator	LABOTERY	GSP-70
Autoclave	PANASONIC	MLS-3751L-PC

1.2 Major reagents

Reagent	Manufacturer	Article Number
Ethanol	SCRC	100092683
BioDewax and Clear solution	Servicebio	G1128
PBS	Servicebio	G0002
Proteinase K	Servicebio	G1234
4% of paraformaldehyde (DEPC water)	Servicebio	G1113
DEPC	Amresco	E174
20×SSC solution	Servicebio	G3016-4
BSA	Servicebio	G5001
DAPI	Servicebio	G1012
Anti-fluorescence quenching sealing tablets	Servicebio	G1401

地址：湖北省武汉市高新区高新二路 388 号生物医药加速器 C22 栋 5 楼 网址：<http://www.servicebio.cn>
电话：4006-027-178

Add: 5th Floor, C22 Building, Biopark, No. 388 Gaoxin 2nd Road, East Lake High-tech Developing Zone,
Wuhan, Hubei, China, 430079 Website: <http://www.servicebio.com> Tel: 027-5111-3188



2. The Steps of the Experiment

2.1 **Organization fixation:** take out the organization, wash clean, then Immediately put in the fixed fluid (DEPC) above 12h.

2.2 **Dehydration:** The tissue is dehydrated by gradient alcohol, paraffin, embedding.

2.3 **Section:** The paraffin is sliced through the slicer, the piece of the slice machine and the 62 - degree oven roast for 2 hours.

2.4 **Dewaxing and dehydration:** Soak sections in 2 changes of Dewaxing Transparent Liquid, 15 minutes each. Dehydrate in 2 changes of pure ethanol for 5 minutes each. Then, followed respectively by dehydrating in gradient ethanol of 85% and 75% ethanol 5 minutes each. Wash in DEPC dilution.

2.5 **Digestion:** according to the tissue fixation time, the slices are boiled in the retrieval solution for 10-15 minutes and naturally cooled. Mark the objective tissue with liquid blocker pen, according to the characteristics of tissues, Add proteinase K (20 ug/ml) working solution to cover objectives and incubate at 37°C for ___ min. Wash in pure water, then wash three times in PBS (pH 7.4) on a Rocker device, 5 min each.

2.6 **Pre-hybridization:** add Pre-hybridization solution to each section and incubate for 1 h at 37°C.

2.7 **Hybridization:** remove the pre-hybridization solution, add the probe hybridization solution with concentration of ___, and incubate the section in a humidity chamber and hybridize overnight at ___°C.

2.8 **Washing:** remove the hybridization solution. Wash sections in 2×SSC for 10 min at 37°C , Wash sections in 1×SSC two times for 5 min each at 37°C, and wash in 0.5×SSC for 10 min at room temperature. Formamide washing can be added if there are more non-specific hybrids.

2.9 **Stain cell nuclei (counter stain):** incubate with DAPI for 8min in the dark, and then mounting.

2.10 **Microscopic examination and photography:** to take photos with positive fluorescence microscope. DAPI glows blue by UV excitation wavelength 330-380 nm and emission wavelength 420 nm; FAM glows green by excitation wavelength 465-495 nm and emission wavelength 515-555 nm; CY3 glows red by excitation wavelength 510-560 nm and emission wavelength 590 nm.

3. Interpretation of the Results

The nuclear stained by DAPI were blue under ultraviolet excitation, and the positive expression



武汉赛维尔生物科技有限公司

Wuhan Servicebio Technology Co., Ltd.

was a kind of fluorescence labeled by corresponding luciferin. FAM (488) is green light, cy3 is red light. The results of mRNA in situ hybridization were cytoplasmic positive and a few nuclear positive were normal. MicRNA and lncRNA were expressed differently. According to the expression, Different fluorescence brightness is strong or weak

Attached table 1 probe information.

地址：湖北省武汉市高新区高新二路 388 号生物医药加速器 C22 栋 5 楼 网址：<http://www.servicebio.cn>
电话：4006-027-178

Add: 5th Floor, C22 Building, Biopark, No. 388 Gaoxin 2nd Road, East Lake High-tech Developing Zone, Wuhan, Hubei, China, 430079 Website: <http://www.servicebio.com> Tel: 027-5111-3188