



细胞荧光 tunel 绿光实验报告

1.实验器材及试剂

1.1 实验器材

名称	厂家	型号
载玻片	Servicebio	
涡旋混合器	Servicebio	MX-F
掌上离心机	Servicebio	D1008E
脱色摇床	Servicebio	TSY-B
移液枪	Dragon	KE0003087/KA0056573
组化笔	Servicebio	G6100
正置荧光显微镜	日本尼康	NIKON ECLIPSE C1
成像系统	日本尼康	NIKON DS-U3

1.2 主要实验试剂

试剂	厂家	货号
PBS 缓冲液	Servicebio	G0002
破膜工作液	Servicebio	G1204
Tunel 试剂盒	Servicebio	G1501
DAPI	Servicebio	G1012
抗荧光淬灭封片剂	Servicebio	G1401

2.细胞爬片荧光 tunel 实验步骤

2.1 细胞破膜：爬片稍甩干后用组化笔在盖玻片中间细胞分布均匀的位置画圈（防止抗体流失），加 50-100 μ l 破膜工作液，室温孵育 20min，PBS 洗 3 次，每次 5 min。

2.2 室温平衡：爬片稍甩干后在圈内滴加 buffer 覆盖组织，buffer 常温孵育 10min。

2.3 加反应液：按爬片数量大小取 tunel 试剂盒内适量 TDT 酶，dUTP，buffer 按 1:5:50 比例混合，加到圈内覆盖细胞，爬片置于 37 $^{\circ}$ C 恒温箱孵育 2 小时。

2.4 DAPI 复染细胞核：爬片用 PBS（PH7.4）洗涤 3 次，每次 5min。去除 PBS 后在圈内滴加 DAPI 染液，避光室温孵育 10min。



武汉赛维尔生物科技有限公司

Wuhan Servicebio Technology Co., Ltd.

2.5 封片：爬片用 PBS (PH7.4) 洗涤 3 次，每次 5min。爬片稍甩干后将细胞的一面朝下用抗荧光淬灭封片剂将玻片封固在载玻片上封片。

2.6 镜检拍照：切片于荧光显微镜下观察并采集图像。（DAPI 紫外激发波长 330-380nm，发射波长 420nm，发蓝光；FITC 激发波长 465-495nm，发射波长 515-555 nm，发绿光；CY3 激发波长 510-560，发射波长 590nm，发红光）。

3.细胞爬片荧光 tunel 结果判读

DAPI 染出来的细胞核在紫外的激发下为蓝色，试剂盒为 FITC 荧光素标记，阳性凋亡细胞核为绿色。

地址：湖北省武汉市高新区高新二路 388 号生物医药加速器 C22 栋 5 楼 网址：<http://www.servicebio.cn>
电话：4006-027-178

Add: 5th Floor, C22 Building, Biopark, No. 388 Gaoxin 2nd Road, East Lake High-tech Developing Zone, Wuhan, Hubei, China, 430079 Website: <http://www.servicebio.com> Tel: 027-5111-3188



Cell (FITC) Tunel Assay Protocol (Cell Climbing Slides)

1. Apparatus and reagents

1.1 Apparatus

Name	Producer	Model
Glass microscope slide	Servicebio	
Vortex	Servicebio	MX-F
Micro-centrifuge	Servicebio	D1008E
Rocker	Servicebio	TSY-B
Pipettor	Dragon	KE0003087/KA0056573
Liquid blocker pen	Servicebio	G6100
Ortho-Fluorescent Microscopy	Nikon	NIKON ECLIPSE C1
Imaging system	Nikon	NIKON DS-U3

1.2 Major reagents

Name	Producer	Model
PBS solution	Servicebio	G0002
Permeabilize solution	Servicebio	G1204
DAPI	Servicebio	G1012
anti-fade mounting medium	Servicebio	G1401
Tunel assay kit	Servicebio	G1501

2. Procedure

2.1 Permeabilization: dry the cell climbing slides slightly. Mark the objective area with liquid blocker pen, where add 50-100 μ l of permeabilize working solution. Incubate for 20 min at room temperature. Wash three times with PBS solution, 5 min each.

2.2 Equilibrium at room temperature: After the slices are slightly dried, buffer is added to the tissues in the circle, and the buffer is incubated at room temperature for 10 minutes

2.3 Tunel reaction: Take appropriate amount of TDT enzyme, dUTP and buffer in the tunel kit according to the number of slices and tissue size and mix at 1:5:50 ratio. Prepare this reaction



solution according to demand before use. Add this mixture to objective tissue placed in a flat wet box, incubate at 37°C for 2 h. be sure to keep the wet box moist by adding water

2.4 DAPI counterstain in nucleus: wash three times with PBS (pH 7.4) in a Rocker device, 5 min each. Then incubate with DAPI solution at room temperature for 10 min, kept in dark place.

2.5 Mount: wash three times with PBS (pH 7.4) in a Rocker device, 5 min each. Throw away liquid slightly, then coverslip with anti-fade mounting medium.

2.6 Microscopic examination and collecting images through fluorescence microscope. DAPI emits blue light at an ultraviolet excitation wavelength of 330-380 nm and an emission wavelength of 420 nm; FITC has an excitation wavelength of 465-495 nm and an emission wavelength of 515-555 nm, and emits green.

3. Results

Nucleus is blue by labeling with DAPI. Tunel assay kit is labelled with FITC. Positive apoptosis cells are green.